

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
25 March 2004 (25.03.2004)

PCT

(10) International Publication Number  
**WO 2004/023879 A1**

(51) International Patent Classification<sup>7</sup>: A21D 2/26,  
8/04, C12N 9/24

la Grotte, 44, B-4130 TILFF (BE). FELLER, Georges  
[BE/BE]; Grosses Pierres, 29, B-4870 TROOZ (BE).

(21) International Application Number:  
PCT/BE2003/000152

(74) Agents: VAN MALDEREN, Joëlle et al.; OFFICE VAN  
MALDEREN, Place Reine Fabiola, 6/1, B-1083 BRUS-  
SELS (BE).

(22) International Filing Date:  
11 September 2003 (11.09.2003)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,  
GH, GM, HN, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,  
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,  
MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,  
RO, RU, SC, SD, SE, SG, SK, SI, SY, TJ, TM, TN, TR,  
TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/410,345 11 September 2002 (11.09.2002) US

(71) Applicant (*for all designated States except US*): PU-  
RATOS NAAMLOZE VENNOOTSCHAP [BE/BE];  
Industrialaan, 25, B-1702 GROOT-BIJGAARDEN (BE).

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): DUTRON, Ag-  
nès [BE/BE]; Boulevard Louis Mettwie, 78, B-1080  
BRUSSELS (BE). GEORIS, Jacques [BE/BE]; Rue  
de la Station, 87, B-6820 FLORENVILLE (BE).  
GENOT, Bernard [BE/BE]; Rue de l'allais, 41, B-4530  
VILLERS-LE-BOUILLET (BE). DAUVIN, Thierry  
[BE/BE]; Rue Bordia, 15, B-4218 COUTHUIN (BE).  
COLLINS, Tony [BE/BE]; Rue des Airelles, 5, B-4031  
ANGLEUR (BE). HOYOUX, Anne [BE/BE]; Avenue de

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: USE OF FAMILY 8 ENZYMES WITH XYLANOLYTIC ACTIVITY IN BAKING

(57) Abstract: The present invention describes a method to improve the properties of a dough and/or a baked product by adding a bread or dough-improving agent containing an enzyme with xylanolytic activity belonging to glycoside hydrolases family 8. Preferred enzymes are the psychrophilic xylanase from *Pseudoalteromonas haloplanktis* and the mesophilic xylanase Y from *Bacillus halodurans* C-125

WO 2004/023879 A1

5

USE OF FAMILY 8 ENZYMES WITH XYLANOLYTIC  
ACTIVITY IN BAKING

Field of the invention

10 [0001] The present invention concerns a method  
and a composition for the improvement of bakery products  
that comprise at least one enzyme with xylanolytic  
activity from the glycoside hydrolase family 8. In an  
embodiment of the invention said enzyme with xylanolytic  
15 activity is further characterized by the fact that there  
is an inversion of configuration during hydrolysis. In  
specific embodiments of the invention said enzyme with  
xylanolytic activity is a psychrophilic xylanase from  
*Pseudoalteromonas haloplanktis* or a mesophilic xylanase  
20 from *Bacillus halodurans*.

Background of the invention

[0002] Xylans are heteropolysaccharides that form  
the major part of the hemicellulose present in the plant  
25 biomass. The backbone of these polysaccharides is a  
chain of  $\beta$  1-4 linked xylopyranosyl residues. Many  
different side groups could bind to these residues like  
acetyl, arabinosyl and glucuronosyl residues. Phenolic  
compounds such as ferulic or hydroxycinnamic acids are  
30 also involved through ester binding in the cross linking  
of the xylan chains or in the linkage between xylan and  
lignin chains for example.

[0003] Endoxylanases (or endo- $\beta$ -1,4-xylanases)  
hydrolyse specifically the backbone of the  
35 hemicellulose. In some cases, the side groups may mask

the main chain by steric hindrance. Different xylanases activities have been described. Their specificity towards their substrate varies from one to each other. Some are more active on insoluble arabinoxylans. The length of the oligomers produced also depends on the type of xylanase considered.

[0004] Glycoside hydrolases (formerly known as cellulase family D) have been classified in 91 families (<http://afmb.cnrs-mrs.fr/CAZY/>) based on sequence homologies, structural and mechanistic features. Because the fold of proteins is better conserved than their sequences, some of the families can be grouped in 'clans' (Henrissat B. 1991, Biochem. J. Vol. 280, p 309). Endo-beta-1,4-xylanases are generally classified in families 10 (formerly family F) and 11 (formerly family G) and are found to frequently have an inverse relationship between their pI and molecular weight. The family 10 xylanases (EXs10) are larger, more complex, while the family 11 xylanases (EXs11) are smaller. Moreover a significant difference in the structure and the catalytic properties of both families exists. EXs10 present an ( $\alpha/\beta$ )<sub>8</sub> barrel fold, have about 40% of  $\alpha$ -helix structures and belong to clan GH-A (Dominguez et al., 1995, Nat. Struct. Biol. Vol. 2, p569) while EXs11 exhibit a  $\beta$ -jelly fold conformation, have about 3-5% of  $\alpha$ -helix structures and belong to clan GH-C (Törrönen et al., 1994, EMBO J. Vol. 13, p2498). EXs10 have a smaller substrate binding site and a lower substrate specificity, have frequently endoglucanase activity and produce smaller oligosaccharides compared to EXs11, which have a higher affinity for unsubstituted xylan (Biely et al., 1997, J. Biotechnol. Vol. 57, p151). All xylanases of both families characterized to date retain the anomeric configuration of the glycosidic oxygen following hydrolysis in which two glutamates function as

the catalytic residues (Jeffries, 1997, Curr. Opin. Biotechnol. Vol. 7, p337).

[0005] Xylanases are used in various industrial areas such as the pulp, paper, feed and bakery industries. Other applications lie in the juice and beer sectors. Xylanases could also be used in the wheat separation process. The observed technological effects are, among others, improved bleachability of the pulp, decreased viscosity of the feed or changes in dough characteristics.

[0006] The use of xylanases (also called hemicellulases or pentosanases) in baking is well known since a number of years. These dough-conditioning enzymes can improve the dough machinability and stability as well as the oven-spring and the crumb structure. Other effects of the enzymes are a larger loaf volume and a softer crumb.

[0007] The mechanism of action of xylanases in bread preparation is still not clearly elucidated. There are about 3 to 4 % pentosans in wheat flour. These pentosans can absorb large amounts of water (up to 30%). This water absorption contributes to the properties of the dough as well as to the quality of the final product. Partial hydrolysis of pentosans by pentosanases into water soluble short chain oligosaccharides increases the water binding capacity. There is also a strong interaction of the pentosans with the gluten fraction of the flour to form a network. Pentosanases may help to relax this strong and rigid network and therefore allowing the carbon dioxide formed by the yeast to better expand the dough.

[0008] Many types of hemicellulase preparations have been used as baking active-ingredients and are commercially available. They are produced by microbial fermentation using various microorganisms as enzyme sources. Some of these enzymes are also produced by

genetically modified microorganisms. All documented uses of xylanases having a positive effect on the specific volume of baked goods, relate to xylanases belonging to glycoside hydrolases Family 10 or Family 11 as defined previously.

[0009] Examples of xylanases for baking are the xylanases from *Bacillus subtilis* and *Aspergillus niger*.

[0010] There is a variety of methods to evaluate the xylanase activity in an enzyme preparation. Examples of such methods of xylanase activity determination are the measure of the release of reducing sugar from xylan (Miller G.L. 1959, Anal. Chem. Vol. 31, p426) or the measure of the release of coloured compounds from modified substrates (for examples AzowAX or Xylazyme AX from Megazyme). However no direct correlation could be found between the xylanolytic activity found in the various enzyme preparations and the effect in baking. Dose-related results can be observed to a certain extend for a single enzyme but the same dosage for two xylanases of different origins does not give the same result in the dough or in the bread. Several reasons could explain this: the differences in substrate specificity, the differences in temperature and pH optimum,...

[0011] It is of great interest to develop new enzyme preparations, such as bread improver compositions or agents, with new or improved properties. One of these properties could be that the xylanase fraction is as small as possible (in terms of weight of enzymes needed to obtain a particular result in baking).

[0012] A xylanase from the Antarctic bacterium *Pseudoalteromonas haloplanktis* has been recently described (Collins, T. et al. 2002. A novel Family 8 xylanase: functional and physico-chemical characterisation. J. Biol. Chem. Vol. 277-38, p35133; Collins, T. et al. 2003, Activity, stability and

flexibility in glycosidases adapted to extreme thermal environments. J. Mol. Biol. Vol. 328, p419; Van Petegem F. et al., 2002. Crystallization and preliminary X-ray analysis of a xylanase from the psychrophile  
5 *Pseudoalteromonas haloplanktis*. Acta Crystallogr D Biol Crystallogr. Vol. 58(Pt 9), p1494-6). This enzyme is a typical psychrophilic enzyme and presents a high catalytic activity at low temperature. It is not homologous to family 10 or 11 xylanases but has 20 to  
10 30% identity with glycoside hydrolases family 8 (formerly family D) members, a family that comprises mainly endoglucanases but also lichenases and chitosanases. Furthermore, a FingerPRINTScan against PRINTS using the InterPro Scan search program (Zdobnov  
15 and Apweiler, 2001, Bioinformatics Vol. 17, p847) indicated that the isolated sequence contained the glycosyl hydrolases family 8 fingerprint and family 8 residues that are strictly conserved in 20 family 8 enzymes analyzed.

20 [0013] Conversely to EXs10 and EXs11, this family 8 xylanase (EXs8) has both high pH and high molecular weight. Structural and catalytic properties are different to those of both EXs10 and EXs11. EXs8 present an ( $\alpha/\alpha$ )<sub>6</sub> barrel fold with 13  $\alpha$ -helices and 13  $\beta$ -strands  
25 and belong to clan GH-M (Van Petegem et al., 2003, The structure of a cold-adapted family 8 xylanase at 1.3 Å resolution. Structural adaptations to cold and investigation of the active site. J. Biol. Chem. Vol. 278(9), p7531-9). These enzymes have no endoglucanase,  
30 chitosanase or licheninase activity and appear to be functionally similar to EXs11, being more active on long chain xylo-oligosaccharides.

[0014] In contrast to other known EXs10 and EXs11 that retain the configuration, family 8 glyco-hydrolases  
35 (Fierobe et al., 1993. Eur. J. Biochem. Vol. 217, p557; <http://afmb.cnrs-mrs.fr/CAZY/>) tend to hydrolyse the

- substrate with inversion of the anomeric configuration of the glycosidic oxygen following hydrolysis in which one glutamate and one aspartate function as the catalytic residues. This has been shown for example for
- 5 the psychrophilic xylanase from *Pseudoalteromonas haloplanktis* (Van Petegem et al., 2003, J. Biol. Chem., Vol. 278(9), p7531-9; Collins, T. et al. 2002. A novel family 8 xylanase: functional and physico-chemical characterization. J. Biol. Chem. Vol. 277(38), p35133).
- 10 Other xylanases belonging to the glycoside hydrolases family 8 have been already described (Yoon, K-H., 1998, Molecular cloning of a *Bacillus* sp. KK-1 xylanase gene and characterization of the gene product. Biochem. Mol. Biol. International. 45(2), p. 337; *Bacillus halodurans*
- 15 C-125 xylanase Y GenBank/GenPept™ accession code BAB05824). These xylanases show sequence homologies between them as well as with the *Pseudoalteromonas haloplanktis* xylanase as described by Collins et al (2002, see above).
- 20 [0015] The list of the enzymes belonging to the glycoside hydrolases family 8 is regularly updated (<http://afmb.cnrs-mrs.fr/CAZY/>).

#### Aims of the invention

- 25 [0016] The present invention aims to provide novel enzyme preparations such as bread improver compositions.
- [0017] The present invention further aims to provide a new method for obtaining improved bakery
- 30 products by using such enzyme preparations.

Summary of the invention

[0018] It was surprisingly found that an enzyme with xylanolytic activity and belonging to glycoside hydrolases family 8 had a positive effect on the dough properties or on the baked product properties when it was added during the mixing of the dough.

[0019] It was also surprisingly found that an enzyme with xylanase activity and hydrolysing with inversion of the anomeric configuration had a positive effect on the dough properties or on the baked products properties when it was added during the mixing of the dough.

[0020] It was further surprisingly found that the use of enzymes as described above increases the volume of the loaf.

[0021] It was also found that the use of some such enzymes with xylanolytic activity from the glycoside hydrolases family 8, the EXs8, make it possible to use less enzyme for the same effect on the dough or bread properties as compared to presently known xylanases.

[0022] A specific embodiment of this invention concerns enzymes with xylanolytic activity from the glycoside hydrolases family 8 and their use in the baking industry as described above.

[0023] A particular embodiment of the invention concerns a psychrophilic family 8 xylanase from *Pseudoalteromonas haloplanktis* and a mesophilic family 8 xylanase Y from *Bacillus halodurans*. These enzymes are xylanases that hydrolyze with inversion of the anomeric configuration.

[0024] Preferred xylanases are those obtained from the strains *Pseudoalteromonas haloplanktis* TAH3a, and *Bacillus halodurans* C-125 or those obtained from the corresponding gene expressed in a suitable host.



[0025] Preferably, the bread improving composition comprising at least one enzyme of the invention is added during the mixing of the dough.

[0026] Said bread improving composition may  
5 further comprise other bread-improving agents selected from the list consisting of other enzymes, emulsifiers, oxidants, milk powder, fats, sugars, amino acids, salts, proteins (gluten, cellulose binding sites) or a mixture thereof.

10 [0027] Said other enzymes may be selected from the list consisting of alpha-amylases, beta-amylases, maltogenic amylases, other xylanases, proteases, glucose oxidase, oxido-reductases, glucanases, cellulases, transglutaminases, isomerases, lipases, phospholipases,  
15 pectinases or a mixture thereof.

[0028] Said alpha-amylase preferably is obtained from *Aspergillus oryzae*.

[0029] The enzymes with xylanolytic activity according to the invention may be used in the form of a  
20 cell extract, a cell-free extract or as a purified protein in said bread improving composition.

[0030] The enzymes of the invention may be mixed with other ingredients and be used in the form of a dry powder or a granulate, in particular a non-dusting  
25 granulate, or in the form of a liquid, preferably with one or more stabilizer(s) such as polyols, sugars, organic acids or sugar alcohols.

[0031] Another aspect of the invention concerns baked products obtained or obtainable when using an  
30 enzyme of the invention.

[0032] A further aspect of the invention concerns bread improving compositions comprising at least one of the enzymes of the invention.

[0033] A still further aspect of the invention  
35 concerns a method for increasing the loaf volume of a

baked product, comprising the step of adding during the mixing of the dough of said baked product, a sufficient amount of an enzyme with xylanolytic activity selected from the group consisting of glycoside hydrolase family 8 xylanases, possibly enzymes with xylanolytic activity that hydrolyze with inversion of configuration. It is possible to use any combination or mixture of the above described enzymes in a dough or bakery product for any of the above described purposes.

10

#### Brief description of the drawings

[0034] The Figure 1 represents the effect of increasing amounts of *Pseudoalteromonas haloplanktis* xylanase TAH3a on the volume of Belgium hard rolls.

15 [0035] The Figure 2 shows a SDS-polyacrylamide gel electrophoresis of various xylanase samples (amount of enzyme loaded proportional to the amount used in baking, same proportionality factor for all enzymes).

Lane 1: molecular weight markers.

20 Lanes 2-5: amounts of *Pseudoalteromonas haloplanktis* xylanase (2-3) and *Bacillus subtilis* xylanase (4-5) needed to treat respectively 0.333 (2, 4) and 0.166 kg (3, 5) of wheat flour.

[0036] The figures 3A and B represent the nucleotide sequence of the xylanase gene of *Pseudoalteromonas haloplanktis* (A) and the corresponding xylanase (B).

[0037] The figure 4 represents the xylanase Y from *Bacillus halodurans* strain C-125.

30

[0038] The invention will be described in further details in the following examples and embodiments by reference to the enclosed drawings. Particular embodiments and examples are not in any way intended to limit the scope of the invention as claimed.

35

**Detailed description of the invention**

[0039] The present invention in particular relates to the use of family 8 enzymes with xylanolytic activity in baking to improve among others the loaf  
5 volume. Said enzymes may have other activities, but the xylanolytic activity should be such that the enzyme is suited for the purposes of the present invention.

**A. Preparation of some preferred enzymes according to**  
10 **the invention**

[0040] Enzymes with xylanolytic activity from the glycosyl hydrolases family 8 can be obtained from various sources (see e.g. <http://afmb.cnrs-mrs.fr/CAZY/> for an overview). Some of the enzymes have been  
15 described in the scientific literature and may therefore be obtained using the techniques described in the corresponding publications. Below, the preparation of some preferred enzymes according to the invention is described in more details.

20

**A.1. Xylanase from *Pseudoalteromonas haloplanktis* TAH3a**

[0041] The xylanase from *Pseudoalteromonas haloplanktis* TAH3a has been described in Collins T. et al. (2002. A novel Family 8 xylanase: functional and  
25 physico-chemical characterisation. J. Biol. Chem. Vol. 277(38), p35133). Collins et al. describes the preparation by liquid culture of a recombinant strain of *Escherichia coli* carrying the *Pseudoalteromonas haloplanktis* TAH3a xylanase gene cloned in an expression  
30 plasmid.

[0042] The nucleotide sequence of the xylanase gene of *Pseudoalteromonas haloplanktis* is available, as well as is available the corresponding amino acid sequence (EMBL Accession number AJ427921, sequence  
35 enclosed by reference thereto, Figures 3A-B, SEQ ID NOs:

6-7). The enzyme has a molecular weight of 46000 Da, an isoelectric point of approximately 9.5, an optimum temperature of 35°C and an optimum pH between 5.3 and 8. It is a true xylanase with the main reaction products of xylan hydrolysis being xylotriase and xyloetraase after prolonged incubation.

[0043] Crude or purified preparation of this xylanase can be obtained by first cultivating either the wild-type of the recombinant strain in a medium suitable to express the enzyme eventually followed by one or several steps of purification such as but not limited to centrifugation, cell disruption, microfiltration, ultrafiltration, precipitation, liquid chromatography, freeze-drying,...

15

#### A.2.Xylanase Y from *Bacillus halodurans* C-125

[0044] The sequence of the family 8 xylanase Y from *Bacillus halodurans* strain C-125 (ATCCCBAA-125) is available as GenBank/GenPept™ accession code BAB05824 (sequence enclosed by reference thereto, Figure 4, SEQ ID NO: 8). The corresponding DNA sequence is available as part as the complete genome sequence of *Bacillus halodurans* strain C-125 (accession code AP001514, sequence enclosed by reference thereto).

[0045] The xylanase from *Bacillus halodurans* C-125 may be obtained by first culturing the strain in/on a medium suitable for expressing the xylanase followed by one or several steps of purification such as but not limited to centrifugation, cell disruption, microfiltration, ultrafiltration, precipitation, liquid chromatography, freeze-drying,...

[0046] Alternatively, the xylanase from *Bacillus halodurans* C-125 may be obtained by first cloning the corresponding gene using well known techniques such as PCR amplification, ligation to a vector and transformation in a microbial host. After this step, the

xylanase may be advantageously expressed in a suitable host by placing the gene under the control of appropriate control sequences such as but not restricted to promoter and terminator and inserting this DNA  
5 construct in a cell that would permit the expression of the gene.

[0047] In a specific embodiment of the present invention the xylanase gene will be overexpressed in said cell.

10

[0048] The present invention is, however, not restricted to the above enzymes but extends to other enzymes with xylanolytic activity belonging to the glycosidase hydrolases family 8. These enzymes are  
15 produced by microbial and non-microbial living cells such as but not restricted to plants or animals. The enzymes may be obtained directly from living cell cultures and/or samples, or may be produced by recombinant strains or host cells. The gene encoding the  
20 enzyme of interest may be a synthetic gene.

[0049] By "recombinant strain" or by "recombinant host cell" is meant a strain having incorporated the nucleotide sequence of said enzymes with xylanolytic activity belonging to the glycosidase hydrolases family  
25 8.

[0050] Advantageously, recombinant host cells are selected from the group consisting of the microbial world, preferably bacteria or fungi, including yeast and more preferably *Bacillus*.

30 [0051] Preferably, said recombinant strain is capable of overexpressing said nucleotide sequence and allows advantageously a high production of the said xylanase.

[0052] The enzymes with xylanolytic activity  
35 produced according to the present invention can be used directly for the purpose of the said present invention

and/or may undergo one or more purification or (further) culturing steps. In a particular embodiment of the invention, the enzymes may be used in pure form.

[0053] Possible preparation methods for enzymes according to the invention include among others preparation or purification from cultures of microorganisms, recombinant or not, in shake flasks or in fermentors, preparation or purification from immobilized cultures, extraction and/or purification from living cells (plants,...),

#### **B. Application of the enzyme**

[0054] The enzyme with xylanolytic activity according to the present invention, purified or not purified, is particularly suited as bread-improving agent. Bread-improving agents are products that could improve or increase texture, flavour, anti-staling effects, softness, crumb softness upon storage, freshness, dough machinability, volume of a final product. Preferably, said enzyme with xylanolytic activity improves the dough handling and/or increases the specific volume of the final baked product. "Baked product" intends to include any product prepared from dough, in particular a yeast raised baked product. Dough is obtained from any type of flour or meal (e.g. based on wheat, rye, barley, oat, or maize). Preferably, dough is prepared with wheat or with mixes including wheat.

[0055] It is one of the aspect of the present invention to show that the enzymes with xylanolytic activity belonging to the family 8 could advantageously be used in a bread improver formula.

[0056] In a further embodiment of the present invention, it is shown that an enzyme with xylanolytic activity belonging to glycoside hydrolases family 8 increases the loaf volume of a baked product.

[0057] In a further embodiment of the present invention, it is shown that the amount of a family 8 enzyme with xylanolytic activity needed to obtain a particular result in baking is exceptionally lower than the amount generally used when working with commercially available enzymes.

[0058] In a further aspect, the present invention relates to the additive effect of said enzyme with xylanolytic activity with other enzymes, in particular with an alpha-amylase, preferably an alpha-amylase from *Aspergillus oryzae*. Said enzyme with xylanolytic activity may be used in combination with other bread-improving agents like but not limited to enzymes, emulsifiers, oxidants, milk powder, fats, sugars, amino acids, salts, proteins (gluten, cellulose binding site), such improving agents being well known to persons skilled in the art. Examples of enzymes are but not restricted to alpha-amylases, beta-amylases, maltogenic amylases, xylanases, proteases, glucose oxidase, oxido-reductases, glucanases, cellulases, transglutaminases, isomerases, lipases, phospholipases, pectinases, etc.

[0059] The enzyme with xylanolytic activity according to the present invention may be used under several forms. Cells expressing the enzyme, such as yeast, fungi, Archea bacteria or bacteria, may be used directly in the process. Said enzyme may be used as a cell extract, a cell-free extract (i.e. portions of the host cell that has been submitted to one or more disruption, centrifugation and/or extraction steps) or as a purified protein. Any of the above-described forms may be used in combination with another enzyme under any of the above-described forms. Said cells, cell extracts, cell-free extracts or enzymes may be mixed with different ingredients, e.g. in the form of a dry powder or a granulate, in particular a non-dusting granulate, in a form of a liquid, for example with stabilisers such

as polyols, sugars, organic acids, sugar alcohols according to established methods.

### Examples

5    Example 1: Preparation of the xylanase of  
     *Pseudoalteromonas haloplanktis* TAH3a

[0060]        The xylanase from *Pseudoalteromonas haloplanktis* was obtained as described (Collins et al, 2002, see above) from a liquid culture of a recombinant  
10    strain of *Escherichia coli* transformed with a DNA vector containing the corresponding gene.

Example 2: Preparation of the xylanase from *Bacillus halodurans* C-125

15    Bacterial strain and culture conditions

[0061]        *Bacillus halodurans* C-125 (ATCC BAA-125) was cultivated in Alkaline Bacillus medium (10g/l glucose , 5g/l yeast extract (Difco), 10g/l Bacto Tryptone (Difco), 1g/l K<sub>2</sub>HPO<sub>4</sub>, 0.2g/l MgSO<sub>4</sub>, 10g/l  
20    Na<sub>2</sub>CO<sub>3</sub>) supplemented with 10g/l birchwood xylan (Sigma) for 72 hours at 37°C. After centrifugation for 1h at 18000g and 4°C the supernatant was concentrated by precipitation with 80% ammonium sulphate and resuspended in 20mM MOPS at pH 8.0. This fraction represents the  
25    crude extract fraction that will be used for the baking trials.

### Cloning and overexpression of the xylanase Y gene

[0062]        Genomic DNA was extracted and purified  
30    from 16 hour cultures, cultivated at 37°C in the modified Alkaline Bacillus medium described above, with the Wizard® Genomic DNA Purification kit (Promega). Based on the published sequence (GenBank/GenPept™ accession AP001514) the entire xylanase Y gene was PCR-



amplified using VENT polymerase (Biolabs Inc, Beverly, MA, USA) with the sense primer

5'-GGGCATATGAAGAAAACGACAGAAGG TG-3', SEQ ID NO: 1

containing an *Nde* I site (underlined)

5 and with the antisense primer

5'-GGCTCGAGCTAG TGTTCCTCTTCTTG-3', SEQ ID NO: 2

containing an *Xho* I site (underlined) and the stop codon (in italics).

[0063] After 3 min initial denaturation at 95 °C, 10 25 cycles of amplification were performed using a Progene apparatus (Techne Cambridge, UK). Each cycle included denaturation at 95 °C for 1 min, hybridization at 52 °C for 30 sec, and elongation at 72 °C for 1.5 min.

15 [0064] The PCR product was cloned into a PCRScript Amp SK(+) vector (Stratagene), using the procedure recommended by the supplier, and used to transform Epicurian Coli XL10-Gold® Kan ultracompetent cells. Blue-white selection allowed selection of white 20 colonies carrying the PCR-fragment. Purified plasmid preparation (Nucleospin plasmid, Macherey-Nagel) was sequenced on an ALF DNA sequencer (Pharmacia Biotech). Sequencing of the inserted fragment was carried out using the universal primers T7 and RP as well as the 25 following primers:

5'-GTGCGGACTGAAGGAATGTC-3', SEQ ID NO: 3

5'-GTATGGTCCATCAACAGAGG-3', SEQ ID NO: 4

5'-GATGGCACTAAAACTCCTGG-3'. SEQ ID NO: 5

[0065] The sequence obtained was identical to the 30 published sequence that could be retrieved from GenBank/genPept™ with the accession number AP001514.

[0066] The xylanase Y gene hence cloned in the PCRScript Amp SK(+) was then excised with *Nde* I and *Xho* I and ligated into the pET 22b(+) cloning vector 35 (Novagen). The resulting recombinant plasmid was transformed to *E. coli* BL21 (DE3) cells (Stratagene).

***Production of the recombinant family 8 xylanase Y from Bacillus halodurans C-125.***

[0067] Fifteen ml of a five hour preculture  
5 (37°C) of the *E. coli* BL21 (DE3) cells carrying the  
xylanase gene was centrifuged at 10000 g for 1 minute  
and the pellet was resuspended in 900 ml Terrific broth  
(12g/l Bacto tryptone (Difco), 24g/l yeast extract  
(Difco), 4ml/l glycerol, 12.54g/l K<sub>2</sub>HPO<sub>4</sub>, 2.31g/l KH<sub>2</sub>PO<sub>4</sub>)  
10 containing 200µg/ml ampicillin in a 3 liter shake flask.  
The culture was incubated at 37°C and 250 rpm until an  
absorbance at 550nm of between 3 - 4 was reached  
whereupon the expression of the enzyme was induced with  
1mM isopropyl-1-thio-β-galactopyranoside. Following 15  
15 hours further incubation at 37°C the cells were  
harvested by centrifugation at 18000 g for 30 minutes at  
4°C, resuspended in 50mM BICINE containing 10mM NaCl,  
disrupted in a prechilled cell disrupter (Constant  
Systems Ltd., Warwick, UK) at 28Kpsi and centrifuged at  
20 40,000 g for 30 minutes.

***Production of the recombinant Bacillus halodurans C-125 family 8 xylanase.***

[0068] The cells of a five hour preculture (37°C)  
25 of the *E. coli* BL21(DE3) cells carrying the xylanase  
gene were collected by centrifugation at 10,000 g for 1  
minute and used to inoculate five liters (15ml  
preculture per liter of culture) of Terrific broth  
(12g/l Bacto tryptone (Difco), 24g/l yeast extract  
(Difco), 4ml/l glycerol, 12.54g/l K<sub>2</sub>HPO<sub>4</sub>, 2.31g/l KH<sub>2</sub>PO<sub>4</sub>)  
30 containing 200µg/ml ampicillin. The culture was  
incubated at 37°C and 250 rpm until an absorbance at  
550nm of between 3 - 4 was reached whereupon the  
expression of the enzyme was induced with 1mM isopropyl-  
35 1-thio-β-galactopyranoside. Following 4 hours further

incubation at 37°C the cells were harvested by centrifugation at 18000 g for 20 minutes at 4°C, resuspended in 20mM MOPS (Sigma), disrupted in a prechilled cell disrupter (Constant Systems Ltd.) at 5 28Kpsi and centrifuged at 40,000 g for 30 minutes. Chromosomal DNA was removed by treatment with 0.2% protamine sulfate (Calbiochem) and centrifugation at 40,000 g for 30 minutes. 25 units of benzonase (Merck, Darmstadt, Germany) was then added and the solution used 10 in baking trials.

**Example 3: Effect of the xylanase of *Pseudoalteromonas haloplanktis* TAH3a in baking of Belgian hard rolls**

[0069] Baking trials were performed to 15 demonstrate the positive effect of *Pseudoalteromonas haloplanktis* TAH3a xylanase in baking. The positive effect was evaluated by the increase in bread volume compared to a reference not containing this enzyme.

[0070] The xylanase was tested in Belgian hard 20 rolls that are produced on a large scale every day in Belgium. The procedure described is well known to the craft baker and it is obvious to one skilled in the art that the same results may be obtained by using equipment from other suppliers.

25 [0071] The ingredients used are listed in the table 1 below:

Table 1

Ingredients (g)	RECIPE 1	RECIPE 2	RECIPE 3	RECIPE 4	RECIPE 5
Flour (Surbi Molens van Deinze)	1500	1500	1500	1500	1500
Water	915	915	915	915	915
Fresh yeast (Bruggeman-Belgium)	90	90	90	90	90
Sodium chloride	30	30	30	30	30
Multec Data HP 20 (Beldem-Belgium)	3.9	3.9	3.9	3.9	3.9
Ascorbic acid	0.12	0.12	0.12	0.12	0.12
Xylanase <i>Pseudoalter. Haloplanktis</i> (units(1)/100kg flour)	0	25	50	100	200

- (1) : One unit of *Pseudoalteromonas haloplanktis* xylanase is defined as the amount of enzyme needed to release 1  $\mu$ mole of reducing sugar (expressed as xylose) from birchwood xylan at 30°C and pH 4.5 (Nelson-Somogyi method)

[0072] The ingredients were mixed for 2 min at low and 8 min at high speed in a Diosna SP24 mixer. The final dough temperature as well as the resting and proofing temperatures were 25°C. After resting for 15 min at 25°C, the dough was reworked manually and rested for another 10 min. Afterwards, 2 kg dough pieces were made up and proofed for 10 min. The 2-kg dough pieces were divided and made up using the Eberhardt Optimat. 66 gr. round dough pieces were obtained. After another 5 min resting time, the dough pieces were cut by pressing and submitted to a final proofing stage for 70 min. The dough pieces were baked at 230°C in a MIWE Condo™ oven with steam (Michael Wenz-Arnstein-Germany). The volume of 6 rolls was measured using the commonly used rapeseed displacement method.

[0073] The results are presented on the table 2 below:

Table 2

Xylanase units	Rolls volume (ml)
0	2325
25	2675
50	2875
100	2950
200	3100

5 [0074] A graphical representation of the effect of the xylanase is shown on figure 1.

[0075] These results show that the xylanase of *Pseudoalteromonas haloplanktis* has a positive effect on the volume of bread.

10

Example 4: Comparison of xylanase amounts used in baking

[0076] The relative amount of xylanases needed to obtain the same effect on the volume in the Belgian hard rolls assay described in Example 3 has been compared by  
15 SDS-polyacrylamide gel electrophoresis.

[0077] The xylanases tested and their respective usage levels in baking were the following:

- the xylanase from *Pseudoalteromonas haloplanktis* TAH3a (30 *P. Haloplanktis* units/100 kg flour)
- 20 - the xylanase from *Bacillus subtilis* (Belase B210 (BELDEM S.A., Belgium) - 3 g/100 kg flour).

[0078] Amounts of enzyme corresponding to the quantity needed to treat 0.333 kg and 0.166 kg of flour, and resulting in the same volume increase, were loaded  
25 on a 12% Tris-HCl precast gel for SDS-polyacrylamide electrophoresis "Ready Gel" (BIO-RAD, Hercules, CA, USA) and run in a "Mini Protean II" (BIO-RAD) apparatus according to the procedure of the manufacturer. Proteins

were stained with Coomassie blue using standard procedure.

[0079] A picture of the stained gel is shown on figure 2.

- 5 [0080] It can be concluded from this picture that the amount of xylanase from *Pseudoalteromonas haloplanktis* TAH3a needed in baking is significantly lower compared to the other.

10 Example 5: Effect of the xylanase of *Pseudoalteromonas haloplanktis* TAH3a in Argentinian breads

- [0081] Baking trials were performed to demonstrate the positive effect of a family 8 xylanases in baking in another recipe than the one used for  
15 example 3, such as for instance in Argentinian breads. The positive effect was evaluated by the increase in bread volume compared to a reference not containing the enzyme and by the width of the cut on the surface of the bread.

- 20 [0082] The enzymes of the invention were tested in Argentinian breads that are typical long bread which need long time proofing (17 hours at 20°C). The procedure described is well known to the craft baker and it is obvious to one skilled in the art that the same  
25 results may be obtained by using equipment from other suppliers.

[0083] The ingredients used are listed in the table 3 below:

30 Table 3

Ingredients (g)	RECIPE 1	RECIPE 2	RECIPE 3	RECIPE 4
Flour (Surbi Molens van Deinze - Belgium)	1500	1500	1500	1500
Water	810	810	810	810
Fresh yeast	5.25	5.25	5.25	5.25

(Bruggeman-Belgium)				
Sodium chloride	30	30	30	30
Standard Improver (1) (Puratos-Belgium)	22.5	22.5	22.5	22.5
Bel'ase B2100 (Beldem - Belgium) (210 units/g)	0	0.15	0	0
Purified Xylanase of <i>Pseudoalteromonas</i> . <i>haloplanktis</i> TAH3a (units(2)/100kg flour)	0	0	25	50

(1) The standard improver used contains: Fungal alpha amylase (Fungamyl 75.000, Novozymes) 1g/100kg flour, vitamin C 10g, Azodicarbonamide 2g/100 kg flour. This is an example of a standard improver. Absolute and  
5 relative amounts of additives can vary according to local adaptation to wheat flour and process

(2) : One unit of xylanase is defined as the amount of enzyme needed to release 1  $\mu$ mole of reducing sugar (expressed as xylose) from birchwood xylan at 30°C and  
10 pH 4.5 (Nelson-Somogyi method)

[0084] The ingredients were mixed for 2 min at low and 7 min at high speed in a Diosna SP24 mixer. The final dough temperature during the resting was 25°C  
15 while it was 20°C during proofing. After resting for 20 min at 25°C, the dough was reworked manually and rested for 17 hours at 20°C during proofing. Afterwards, 2 kg dough pieces were made up and proofed for 10 min. The 0.35-kg dough pieces were divided and made up using the  
20 Bertrand R8/L8 (+/-30 cm) and submitted to a final proofing stage for 17 hours. The dough pieces were baked at 210°C for 30 min in a MIWE Condo™ oven with steam (Michael Wenz-Arnstein-Germany). The volume of Argentinian breads was measured using the commonly used  
25 rapeseed displacement method.

[0085] The results are presented on the table 4 below:

Table 4

Recipe	Sample	Xylanase units	Bread volume (ml)	Width of cut (mm)
1		0	1675	19
2	Bel'ase B210	2100	1900	27
3	TAH3a	25	1900	34
4	TAH3a	50	1925	30

5 [0086] These results show that the family 8 xylanase has a positive effect on bread making with long time proofing at lower temperature (i.e., 20°C). The amount of the commercial enzyme (Bel'ase B210, see above) used here is 3-fold higher than the optimal dose required as described in example 3 for Belgian hard rolls, while the amount of family 8 xylanase needed remained unchanged. There exists also a difference between the enzyme concentration needed to optimize the width of the cut and the one needed for an optimal volume increase.

Example 6: The effect of various enzymes of the invention versus other enzymes in baking

20 [0087] Baking trials were performed to demonstrate the positive effect of various enzymes with xylanolytic activity belonging to the glycoside hydrolases family 8 in baking. The positive effect was evaluated by the increase in bread volume compared to a reference not containing these enzymes as well as to a commercial enzyme preparation (Bel'ase B210, see above).

25 [0088] The xylanase from *Pseudoalteromonas haloplanktis* TAH3a was prepared and purified as described in example 1. The xylanase Y from *Bacillus*



*halodurans* C-125 was prepared from a recombinant *Escherichia coli* strain as described in example 2.

[0089] The method used to evaluate the performances of the two enzymes was a mini baking test consisting of preparing dough with 100g of flour.

[0090] The procedure described is well known to one skilled in the art and it is obvious that the same results may be obtained by using equipment from other suppliers.

10 [0091] The ingredients used are listed in the table 5 below:

**Table 5**

Ingredients (g)	RECIPE 1	RECIPE 2	RECIPE 3	RECIPE 4	RECIPE 5	RECIPE 6	RECIPE 7
Flour (Surbi -Molens van Deinze)	100	100	100	100	100	100	100
Water	57	57	57	57	57	57	57
Fresh yeast (Bruggeman-Belgium)	5	5	5	5	5	5	5
Sodium chloride	2	2	2	2	2	2	2
Dextrose	2	2	2	2	2	2	2
Ascorbic acid (g/100kg flour)	4	4	4	4	4	4	4
Bel'ase B210 (Beldem - Belgium) (units(1)/100 kg flour)	0	630	1050	0	0	0	0
Purified xylanase from <i>Pseudoalteromonas</i> <i>haloplanktis</i> TAH3a (units(1)/100kg flour)	0	0	0	25	50	0	0
Recombinant xylanase from <i>Bacillus</i> <i>halodurans</i> C-125 (units(1)/100kg flour)	0	0	0	0	0	6000	12000

(1) : One unit of xylanase is defined as the amount of  
15 enzyme needed to release 1  $\mu$ mole of reducing sugar

(expressed as xylose) from birchwood xylan at 30°C and pH 4.5 (Nelson-Somogyi method)

[0092] The ingredients were mixed for 4.5 min in a National mixer. 150 g dough pieces were weighted and rested for 20 min at 25°C in plastic boxes. The doughs were reworked and rested for another 20 min. The final proofing time was 50 min at 36°C. The dough pieces were then baked at 225°C for 20 min. The volume of the breads was measured using the commonly used rapeseed displacement method.

[0093] The results are presented on the table 6 below:

15 **Table 6**

Recipe	Sample	Xylanase units/ 100kg flour	Bread volume (ml)
1		0	650
2	Bel'ase B210	630	680
3	Bel'ase B210	1050	705
4	TAH3a	25	690
5	TAH3a	50	760
6	C-125	6000	725
7	C-125	12000	725

[0094] These results show that the positive effect of enzymes of the invention is not restricted to the *Pseudoalteromonas haloplanktis* TAH3a xylanase. Also other enzyme preparations with xylanolytic activity belonging to the glycoside hydrolase family 8 could improve the properties of bread (demonstrated here for the *Bacillus halodurans* C-125 xylanase, other results not shown).

CLAIMS

1. Method for the preparation of a baked product, said method comprising the step of adding to a  
5 dough of said baked product a bread improving composition comprising at least one enzyme with xylanolytic activity selected from the group consisting of glycoside hydrolases Family 8.

2. The method of claim 1, wherein said  
10 enzyme hydrolyses with inversion of configuration.

3. The method of claim 1 or 2, wherein that said enzyme is a xylanase obtained from the strain *Pseudoalteromonas haloplanktis*.

4. The method of claim 3, wherein said  
15 *Pseudoalteromonas haloplanktis* strain is *Pseudoalteromonas haloplanktis* TAH3a.

5. The method of claim 1 or 2, wherein that said enzyme is a xylanase obtained from the strain *Bacillus halodurans*.

20 6. The method of claim 5, wherein said *Bacillus halodurans* strain is *Bacillus halodurans* C-125.

7. The method of anyone of the preceding claims, wherein said bread improving composition is added during the mixing of the dough.

25 8. The method of anyone of the preceding claims, wherein said bread improving composition further comprises another bread-improving agent which is selected from the list consisting of other enzymes, emulsifiers, oxidants, milk powder, fats, sugars, amino  
30 acids, salts, proteins (gluten, cellulose binding sites) or a mixture thereof.

9. The method of claim 8, wherein said  
other enzyme is selected from the list consisting of alpha-amylases, beta-amylases, maltogenic amylases,  
35 other xylanases, proteases, glucose oxidase, oxido-

reductases, glucanases, cellulases, transglutaminases, isomerases, lipases, phospholipases, pectinases or a mixture thereof.

10. The method of claim 9, wherein said  
5 alpha-amylase is an alpha-amylase obtained from *Aspergillus oryzae*.

11. The method of anyone of the preceding claims, wherein said enzyme with xylanolytic activity is present as a cell extract, a cell-free extract or as a  
10 purified protein.

12. The method of anyone of the preceding claims, wherein said enzyme with xylanolytic activity is mixed with other ingredients in the form of a dry powder or a granulate, in particular a non-dusting granulate,  
15 or in the form of a liquid, preferably with one or more stabilizer(s) such as polyols, sugars, organic acids or sugar alcohols.

13. Baked product obtainable by the method of anyone of the preceding claims.

20 14. The baked product of claim 13, which is an Argentinian bread with an increased width of cut.

15. Method for increasing the loaf volume of a baked product, comprising the step of adding during the mixing of the dough of said baked product, a  
25 sufficient amount of an enzyme with xylanolytic activity selected from the group consisting of glycoside hydrolase family 8 xylanases.

16. Method for increasing the loaf volume of a baked product or for increasing the width of cut on  
30 the surface of a baked product, comprising the step of adding during the mixing of the dough of said baked product, a sufficient amount an enzyme with xylanolytic activity selected from the group consisting of glycoside hydrolase family 8 xylanases, of a family 8 glycoside  
35 hydrolase that hydrolyzes with inversion of configuration and/or of a combination thereof.

17. Bread improver composition for increasing the loaf volume of a baked product or for increasing the width of cut on the surface of a baked product, characterised in that it comprises at least one  
5 enzyme with xylanolytic activity selected from the group consisting of glycoside hydrolases Family 8.

18. The bread improver composition of claim 17, wherein said enzyme hydrolyses with inversion of configuration.



10 19. The bread improver composition of claim 17 or 18, wherein that said enzyme is a xylanase obtained from the strain *Pseudoalteromonas haloplanktis*.

20. The bread improver composition of claim 19, wherein said *Pseudoalteromonas haloplanktis*  
15 strain is *Pseudoalteromonas haloplanktis* TAH3a.

21. The bread improver composition of claim 17 or 18, wherein that said enzyme is a xylanase obtained from the strain *Bacillus halodurans*.

22. The bread improver composition of  
20 claim 21, wherein said *Bacillus halodurans* strain is *Bacillus halodurans* C-125.

1/3

**Fig. 1****Fig. 2**

BEST AVAILABLE COPY

2/3

```

atgaaagtat tttttaaaat aacaacttta ttgttaatac taataagcta
tcaatcactt gctgcattta ataataaccc atcgagtgtg ggcgcctaca
gttcagggac ataccgtaac ctgcgacaag aaatgggtaa aacaaatata
cagcaaaaagg tgaatagtag ttttgacaat atgtttggct ataacaacac
acaacaactt tactaccgtt acaccgaaaa cgggtgtttat aaagcacact
acataaaaagc aattaaccca gacgaaggcg acgatataag aacagaaggg
caatcgtggg gaatgaccgc cgctgtcatg cttataaac aagaagaatt
tgataaccta tggcgctttg caaaagcgta tcaaaaaaat ccagacaatc
accctgatgc taaaaaaciaa ggcgtttacg cgtggaaact aaagcttaat
caaaacggct ttgtttataa agtggatgag ggccccgctc ccgatggcga
agagtacttt gcgtttgcac tacttaatgc ctctgctcgt tgggggaatt
cgggtgagtt taactactac aacgatgcca ttaccatgtt aaacacaatt
aaaaataagc tgatggaaaa ccaaataatc cgcttttcac cttacattga
taacctaaca gacccttctt accatatacc tgcgttttac gactactttg
caaataacgt aactaaccaa gcagacaaaa attactggcg acaagtagcc
acaaaaagta gaaccttact taaaaacat tttacaaaag taagtggtag
cccgcatggg aacttaccta catTTTTatc gcgcttagat ggcagccctg
ttattggcta catTTTTaac ggccaagcaa acccaggtca atggtatgaa
tttgatgcat ggcgcgtaat tatgaatgtg ggttttagacg cgcatttaat
gggtgctcaa gcgtggcata aaagtgcagt taataaagca ctgggctttt
taagttatgc aaaaacaaac aacagtaaaa actgttacga gcaagtgtat
tcgtacgggtg gagcgcaaaa cagaggctgt gcaggcgaag gtcaaaaagc
cgcgaaatgca gtagcgttac ttgcttcaac aaatgctggg caagcaaatg
agttttttta cgaatttttg tctttatcgc aaccaacggg tgactaccgt
tactataatg gttcgttata tatgttagct atgctgcatg tatcgggcaa
ttttaagttt tataacaaca cgtttaatta a (SEQ ID NO: 6)

```

(EMBL Accession number AJ427921)

**Fig. 3A**

```

MKVFFKITLLLLILISYQSLAAFNNNPSSVGAYSSGTYRNLAQEMGKTNIQQKVNSTFD
NMFQYNNNTQQLYYPYTENGVIKAHYIKAINPDEGDDIRTEGQSWGMTAAVMLNKQEEFD
NLWRFKAYQKNPDNHPDAKKQGVYAWKLKLNQNGFVYKVDEGPAPDGEEYFADFALLNA
SARWGNSEGFNYNDAITMLNTIKNKLMEQIIRFSPYIDNLTDPYHIIPAFYDYFANN
VTNQADKNYWRQVATKSRLLKNHFTKVSQSPHWNLPFLSRLDGSPVIGYIFNGQANP
GQWYEFDAWRVIMNVGLDAHLMGAQAWHKS AVNKA LGFLSYAKTNN SKNCYEQVSYGG
AQNRGCAGEGQKAANAVALLASTNAGQANEFFNEFWLSQPTGDYRYNGSLYMLAMLH
VSGNFKFYNNTFN (SEQ ID NO: 7)

```

(EMBL Accession number AJ427921)

**Fig. 3B**

BEST AVAILABLE COPY

MKKTTEGAFYTREYRNLFKEFGYSEAEIQERVKDTWEQLFGDNPETKIYYEVGDDLGYL  
LDTGNLDVRTEGMSYGMMMAVQMDRKDIFDRIWNWTMKNMYMTEGVHAGYFAWSCQPDG  
TKNSWGPAPDGEEYFALALFFASHRWGDGDEQPFNYSEQARKLLHTCVHNGEGGPGHPM  
WNRDNKLIKFIPEVEFSDPSYHLPHFYELFSLWANEEDRVFWKEAAEASREYLKIACHP  
ETGLAPEYAYYDGTNPDEKGYGHFFSDSYRVAANIGLDAEWFGGSEWSAEEINKIQAFF  
ADKEPEDYRRYKIDGEPFEEKSLHPVGLIATNAMGSLASVDGPYAKANVDLFWNTPVVRT  
GNRRYYDNCLYLFAMLALSGNFKIWFPEGQEEH (SEQ ID NO: 8)

(GenBank/GenPept™ accession code BAB05824)

**Fig. 4**

**BEST AVAILABLE COPY**



## SEQUENCE LISTING

<110> PURATOS N.V.  
<120> USE OF FAMILY 8 ENZYMES WITH XYLANOLYTIC ACTIVITY IN BAKING  
<130> P.PURA.19/WO  
<160> 8  
<170> PatentIn version 3.2  
<210> 1  
<211> 28  
<212> DNA  
<213> Artificial  
<220>  
<223> sense primer  
<400> 1  
gggcatatga agaaaacgac agaaggtg 28  
  
<210> 2  
<211> 26  
<212> DNA  
<213> Artificial  
<220>  
<223> antisense primer  
<400> 2  
ggctcgagct agtgttcctc ttcttg 26  
  
<210> 3  
<211> 20  
<212> DNA  
<213> Artificial  
<220>  
<223> sequencing primer  
<400> 3  
gtgctggactg aaggaatgtc 20  
  
<210> 4  
<211> 20  
<212> DNA  
<213> Artificial  
<220>  
<223> sequencing primer  
<400> 4  
gtatggtcca tcaacagagg 20  
  
<210> 5  
<211> 21  
<212> DNA  
<213> Artificial  
<220>  
<223> sequencing primer

<400> 5  
gatggcacta aaaactcctg g 21

<210> 6  
<211> 1281  
<212> DNA  
<213> Pseudoalteromonas haloplanktis

<220>  
<221> misc\_feature  
<223> xylanase gene of Pseudoalteromonas haloplanktis (EMBL accession number AJ427921)

<400> 6  
atgaaagtat tttttaaaat aacaacttta ttgttaatac taataagcta tcaatcactt 60  
gctgcattta ataataaccc atcgagtgtg ggcgcctaca gttcagggac ataccgtaac 120  
ctcgcacaaag aaatgggtaa acaaatata cagcaaaagg tgaatagtac ttttgacaat 180  
atgtttggct ataacaacac acaacaactt tactacccgt acaccgaaaa cgggtgtttat 240  
aaagcacact acataaaagc aattaaccca gacgaaggcg acgatataag aacagaaggg 300  
caatcgtggg gaatgaccgc cgctgtcatg cttataaac aagaagaatt tgataaccta 360  
tggcgctttg caaaagcgta tcaaaaaaat ccagacaatc accctgatgc taaaaaacia 420  
ggcgtttacg cgtggaaact aaagcttaat caaaacggct ttgtttataa agtggatgag 480  
ggccccgctc ccgatggcga agagtacttt gcgtttgcac tacttaatgc ctctgctcgt 540  
tgggggaatt cgggtgagtt taactactac aacgatgcca ttaccatggt aaacacaatt 600  
aaaaataagc tgatggaaaa ccaaataatc cgcttttcac cttacattga taacctaaca 660  
gacccctctt accatatacc tgcgttttac gactactttg caaataacgt aactaaccaa 720  
gcagacaaaa attactggcg acaagtagcc acaaaaagta gaaccttact taaaaaccat 780  
tttacaaaag taagtggtag cccgcattgg aacttaccta cttttttatc gcgcttagat 840  
ggcagccctg ttattggcta cttttttaac ggccaagcaa acccaggtca atggtatgaa 900  
tttgatgcat ggcgcgtaat tatgaatgtg ggttttagacg cgcatttaat ggggtgctcaa 960  
gcgtggcata aaagtgcagt taataaagca ctgggctttt taagttatgc aaaaacaaac 1020  
aacagtaaaa actgttacga gcaagtgtat tcgtacggtg gagcgcaaaa cagaggctgt 1080  
gcaggcgaag gtcaaaaagc cgcgaatgca gtagcgttac ttgcttcaac aaatgctggg 1140  
caagcaaatg agttttttta cgaatttttg tctttatcgc aaccaacggg tgactaccgt 1200  
tactataatg gttcgttata tatgttagct atgctgcatg tatcgggcaa ttttaagttt 1260  
tataacaaca cgtttaatta a 1281

<210> 7  
<211> 426  
<212> PRT  
<213> Pseudoalteromonas haloplanktis

<220>

&lt;221&gt; misc\_feature

<223> xylanase of *Pseudoalteromonas haloplanktis* (EMBL accession number AJ427921)

&lt;400&gt; 7

Met Lys Val Phe Phe Lys Ile Thr Thr Leu Leu Leu Ile Leu Ile Ser  
1 5 10 15Tyr Gln Ser Leu Ala Ala Phe Asn Asn Asn Pro Ser Ser Val Gly Ala  
20 25 30Tyr Ser Ser Gly Thr Tyr Arg Asn Leu Ala Gln Glu Met Gly Lys Thr  
35 40 45Asn Ile Gln Gln Lys Val Asn Ser Thr Phe Asp Asn Met Phe Gly Tyr  
50 55 60Asn Asn Thr Gln Gln Leu Tyr Tyr Pro Tyr Thr Glu Asn Gly Val Tyr  
65 70 75 80Lys Ala His Tyr Ile Lys Ala Ile Asn Pro Asp Glu Gly Asp Asp Ile  
85 90 95Arg Thr Glu Gly Gln Ser Trp Gly Met Thr Ala Ala Val Met Leu Asn  
100 105 110Lys Gln Glu Glu Phe Asp Asn Leu Trp Arg Phe Ala Lys Ala Tyr Gln  
115 120 125Lys Asn Pro Asp Asn His Pro Asp Ala Lys Lys Gln Gly Val Tyr Ala  
130 135 140Trp Lys Leu Lys Leu Asn Gln Asn Gly Phe Val Tyr Lys Val Asp Glu  
145 150 155 160Gly Pro Ala Pro Asp Gly Glu Glu Tyr Phe Ala Phe Ala Leu Leu Asn  
165 170 175Ala Ser Ala Arg Trp Gly Asn Ser Gly Glu Phe Asn Tyr Tyr Asn Asp  
180 185 190Ala Ile Thr Met Leu Asn Thr Ile Lys Asn Lys Leu Met Glu Asn Gln  
195 200 205Ile Ile Arg Phe Ser Pro Tyr Ile Asp Asn Leu Thr Asp Pro Ser Tyr  
210 215 220His Ile Pro Ala Phe Tyr Asp Tyr Phe Ala Asn Asn Val Thr Asn Gln  
225 230 235 240Ala Asp Lys Asn Tyr Trp Arg Gln Val Ala Thr Lys Ser Arg Thr Leu  
245 250 255

Leu Lys Asn His Phe Thr Lys Val Ser Gly Ser Pro His Trp Asn Leu  
 260 265 270

Pro Thr Phe Leu Ser Arg Leu Asp Gly Ser Pro Val Ile Gly Tyr Ile  
 275 280 285

Phe Asn Gly Gln Ala Asn Pro Gly Gln Trp Tyr Glu Phe Asp Ala Trp  
 290 295 300

Arg Val Ile Met Asn Val Gly Leu Asp Ala His Leu Met Gly Ala Gln  
 305 310 315 320

Ala Trp His Lys Ser Ala Val Asn Lys Ala Leu Gly Phe Leu Ser Tyr  
 325 330 335

Ala Lys Thr Asn Asn Ser Lys Asn Cys Tyr Glu Gln Val Tyr Ser Tyr  
 340 345 350

Gly Gly Ala Gln Asn Arg Gly Cys Ala Gly Glu Gly Gln Lys Ala Ala  
 355 360 365

Asn Ala Val Ala Leu Leu Ala Ser Thr Asn Ala Gly Gln Ala Asn Glu  
 370 375 380

Phe Phe Asn Glu Phe Trp Ser Leu Ser Gln Pro Thr Gly Asp Tyr Arg  
 385 390 395 400

Tyr Tyr Asn Gly Ser Leu Tyr Met Leu Ala Met Leu His Val Ser Gly  
 405 410 415

Asn Phe Lys Phe Tyr Asn Asn Thr Phe Asn  
 420 425

<210> 8  
 <211> 388  
 <212> PRT  
 <213> Bacillus halodurans

<220>  
 <221> misc\_feature  
 <223> xylanase Y from Bacillus halodurans strain C-125  
 (GenBank/GenPept™ accession code BAB05824)

<400> 8

Met Lys Lys Thr Thr Glu Gly Ala Phe Tyr Thr Arg Glu Tyr Arg Asn  
 1 5 10 15

Leu Phe Lys Glu Phe Gly Tyr Ser Glu Ala Glu Ile Gln Glu Arg Val  
 20 25 30

Lys Asp Thr Trp Glu Gln Leu Phe Gly Asp Asn Pro Glu Thr Lys Ile  
 Page 4

35                      40                      45  
 Tyr Tyr Glu Val Gly Asp Asp Leu Gly Tyr Leu Leu Asp Thr Gly Asn  
     50                      55                      60  
 Leu Asp Val Arg Thr Glu Gly Met Ser Tyr Gly Met Met Met Ala Val  
     65                      70                      75                      80  
 Gln Met Asp Arg Lys Asp Ile Phe Asp Arg Ile Trp Asn Trp Thr Met  
                     85                      90                      95  
 Lys Asn Met Tyr Met Thr Glu Gly Val His Ala Gly Tyr Phe Ala Trp  
                     100                      105                      110  
 Ser Cys Gln Pro Asp Gly Thr Lys Asn Ser Trp Gly Pro Ala Pro Asp  
                     115                      120                      125  
 Gly Glu Glu Tyr Phe Ala Leu Ala Leu Phe Phe Ala Ser His Arg Trp  
     130                      135                      140  
 Gly Asp Gly Asp Glu Gln Pro Phe Asn Tyr Ser Glu Gln Ala Arg Lys  
     145                      150                      155                      160  
 Leu Leu His Thr Cys Val His Asn Gly Glu Gly Gly Pro Gly His Pro  
                     165                      170                      175  
 Met Trp Asn Arg Asp Asn Lys Leu Ile Lys Phe Ile Pro Glu Val Glu  
                     180                      185                      190  
 Phe Ser Asp Pro Ser Tyr His Leu Pro His Phe Tyr Glu Leu Phe Ser  
                     195                      200                      205  
 Leu Trp Ala Asn Glu Glu Asp Arg Val Phe Trp Lys Glu Ala Ala Glu  
     210                      215                      220  
 Ala Ser Arg Glu Tyr Leu Lys Ile Ala Cys His Pro Glu Thr Gly Leu  
     225                      230                      235                      240  
 Ala Pro Glu Tyr Ala Tyr Tyr Asp Gly Thr Pro Asn Asp Glu Lys Gly  
                     245                      250                      255  
 Tyr Gly His Phe Phe Ser Asp Ser Tyr Arg Val Ala Ala Asn Ile Gly  
                     260                      265                      270  
 Leu Asp Ala Glu Trp Phe Gly Gly Ser Glu Trp Ser Ala Glu Glu Ile  
                     275                      280                      285  
 Asn Lys Ile Gln Ala Phe Phe Ala Asp Lys Glu Pro Glu Asp Tyr Arg  
     290                      295                      300  
 Arg Tyr Lys Ile Asp Gly Glu Pro Phe Glu Glu Lys Ser Leu His Pro

Page 6

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/BE 03/00152

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A21D2/26 A21D8/04 C12N9/24

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A21D C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	COLLINS ET. AL: "A Novel family 8 xylanase: functional and physico-chemical characterisation" J. BIOL. CHEM, vol. 277, no. 38, 8 May 2002 (2002-05-08) - 27 June 2002 (2002-06-27), pages 35133-35139, XP001156740 USA cited in the application the whole document ---	1-22
X	WO 01 70998 A (TEUFEL DANIEL PAUL ;DARAN JEAN MARC GEORGES (FR); HERWEIJER MARGAR) 27 September 2001 (2001-09-27) page 10, paragraph B -page 11 page 28, paragraph 4 -page 29, paragraph 2; claims 1-37 --- -/--	1-22



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

10 December 2003

Date of mailing of the international search report

05/01/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Adechy, M

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/BE 03/00152

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 02 19828 A (NOVOZYMES AS) 14 March 2002 (2002-03-14) page 4, line 24 -page 9, line 17 page 10, line 27 -page 11, line 5; claims 1-38; examples 1-3 ----	1-22
P,A	US 2003/082779 A1 (GEODEGEBUUR FRITS ET AL) 1 May 2003 (2003-05-01) paragraphs '0080!, '0081!, '0127!, '0145! ----	
P,A	US 2003/059496 A1 (CONTRERAS ROBERTO ET AL) 27 March 2003 (2003-03-27) the whole document -----	



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/BE 03/00152

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0170998	A	27-09-2001	AU 6211601 A	03-10-2001
			CA 2403486 A1	27-09-2001
			CN 1426470 T	25-06-2003
			WO 0170998 A1	27-09-2001
			EP 1272643 A1	08-01-2003
			NO 20024516 A	20-11-2002
WO 0219828	A	14-03-2002	AU 8381701 A	22-03-2002
			WO 0219828 A1	14-03-2002
			US 2002094367 A1	18-07-2002
US 2003082779	A1	01-05-2003	WO 03027306 A2	03-04-2003
US 2003059496	A1	27-03-2003	US 2002102326 A1	01-08-2002
			GB 2371205 A	24-07-2002